

MANUFACTURING MILK LAW OF 2001 (EXCERPT)

Act 267 of 2001

ARTICLE 7

288.630 Prohibited acts; standards; compliance.

Sec. 70. (1) A person shall not directly, through an agent, or on behalf of another person sell or offer for sale, furnish, or possess or control with intent to sell or offer for sale, or furnish an unsanitary, adulterated, or misbranded milk or dairy product to a person or a processor.

(2) Dairy products made or sold in Michigan shall comply with the requirements of this act and the standards as follows:

Chemical, Physical, Bacteriological, and Temperature Standards		
MANUFACTURING GRADE RAW MILK FOR PASTEURIZATION (NOT FOR FROZEN DESSERTS) INCLUDING ULTRAFILTRATION OR REVERSE OSMOSIS RAW MILK CONCENTRATE	Temperature	Bulk milk cooled to 45°F (7°C) or less within 2 hours after milking and maintained thereat. Provided that the blend temperature after the first and subsequent milkings does not exceed 50°F (10°C). Can milk not to exceed 60°F (16°C) if used for cheese making; if delivered to the plant within 2 hours of milking, no temperature limit.
	Bacterial limits	Not to exceed 500,000 per ml (milk for cheese not to exceed 750,000 per ml). Not to exceed 1,000,000 per ml as commingled milk prior to pasteurization.
	Somatic cell count	Not to exceed 1,000,000 per ml.
	Drug residues	No positive results on drug residue detection methods which have been found to be acceptable for use with raw milk.
	Sediment	Not to exceed a USDA no. 3 standard following procedures from standard methods for the examination of milk and milk products.
	Freezing point	-0.530°H maximum.
RAW MILK FOR FROZEN DESSERTS	Temperature	Bulk milk cooled to 45°F (7°C) or less within 2 hours after milking and maintained thereat. Provided, that the blend temperature after the first and subsequent

		milking does not exceed 50°F (10°C).
	Bacterial limits	Not to exceed 100,000 per ml for individual supplies, not to exceed 300,000 per ml commingled.
	Somatic cell counts	Not to exceed 750,000 per ml.
	Drug residues	No positive results on drug residue detection methods which have been found to be acceptable for use with raw milk.
	Sediment	Not to exceed a USDA no. 3 standard following procedures from standard methods for the examination of milk and milk products.
	Freezing point	-0.530°H maximum.
PASTEURIZED CONDENSED MILK AND CONDENSED SKIM MILK	Temperature	Cooled to 45°F (7°C) [50°F (10°C) if 45% or more solids] or less, or heated to 145° (63°C) or greater and maintained thereat unless the product is being dried within 4 hours after condensing.
	Bacterial limits	Not to exceed 30,000 per gram.
	Coliform count	Not to exceed 10 per gram. Provided, that in the case of bulk milk transport tank shipments shall not exceed 100 per ml.
	Drug residues	No positive results on drug residue detection methods which have been found to be acceptable for use with condensed milk and condensed skim milk.
<hr/>		
DRY WHOLE MILK, EXTRA GRADE	No more than:	
	Butterfat	Not less than 26% or more than 40%.

Moisture	4.50%.
Solubility index	1.0 ml spray process; 15.0 roller process.
Bacterial limit	Not to exceed 50,000 per gram.
Coliform count	Not to exceed 10 per gram.
Scorched particles disc B	15.0/gram spray process; 22.5 roller process.
Drug residues	No positive results on drug residue detection methods which have been found to be acceptable for use with dry whole milk.
DMCC count	Less than 100,000,000 per gram.

DRY WHOLE MILK,
STANDARD
GRADE

No more than:

Butterfat	Not less than 26% or more than 40%.
Moisture	5.00%.
Titratable acidity	0.15%.
Solubility index	1.5 ml spray process; 15.0 ml roller process.
Bacterial limit	Not to exceed 100,000 per gram.
Coliform count	Not to exceed 10 per gram.
Scorched particles disc B	22.5 per gram spray process; 32.5 per gram roller process.
Drug residues	No positive results on drug residue detection methods which have been found to be acceptable for use with dry, whole milk.
DMCC count	Less than 100,000,000 per gram.

NONFAT DRY MILK, EXTRA

Rendered Friday, February 17, 2017

Page 3

Michigan Compiled Laws Complete Through PA 471 of 2016

© Legislative Council, State of Michigan

Courtesy of www.legislature.mi.gov

GRADE	No more than:	
	Butterfat	1.25%.
	Moisture	4.00%.
	Titratable acidity	0.15%.
	Solubility index	1.2 ml (2.0 ml high-heat, max) spray process; 15.0 ml roller process.
	Bacterial limit	Not to exceed 10,000 per gram spray or 50,000 per gram roller process.
	Coliform count	Not to exceed 10 per gram.
	Scorched particles disc B	15.0/gram spray; 22.5/gram roller process.
	Drug residues	No positive results on drug residue detection methods which have been found to be acceptable for use with nonfat dry milk.
	DMCC count	Less than 100,000,000 per gram.

NONFAT DRY MILK,
STANDARD GRADE

No more than:	
Butterfat	1.50%.
Moisture	5.00%.
Titratable acidity	0.17%.
Solubility index	2.5 ml spray process; 15.0 ml roller process.
Bacterial estimate	75,000/gram spray; 100,000/gram roller process.
Coliform count	10 per gram.
Scorched particles disc B	22.5/gram spray; 32.5/gram roller process.
Drug residues	No positive results on drug residue detection methods which have been found to be acceptable

		for use with nonfat dry milk.
	DMCC count	Less than 100,000,000 per gram.
<hr/>		
INSTANT NONFAT DRY MILK, EXTRA GRADE	No more than:	
	Butterfat	1.25%.
	Moisture	4.50%.
	Titratable acidity	0.15%.
	Solubility index	1.0 ml.
	Bacterial limit	Not to exceed 10,000 per gram.
	Coliform count	Not to exceed 10 per gram.
	Scorched particles disc B	15.0/gram.
	Drug residues	No positive results on drug residue detection methods which have been found to be acceptable for use with nonfat dry milk.
	Dispersibility	85.0%.
	DMCC count	Less than 40,000,000 per gram.
<hr/>		
WHEY FOR CONDENSING	Temperature	Maintained at a temperature of 45°F (7°C) or less, or 135°F (57°C) or greater, except for acid-type whey with a titratable acidity 0.40% or above, or a pH of 4.6 or below.
	Drug residues	No positive results on drug residue detection methods which have been found to be acceptable for use with whey.
<hr/>		
PASTEURIZED CONDENSED WHEY AND WHEY PRODUCTS	Temperature	Cooled to 50°F (10°C) or less during crystallization, within 72 hours of

		condensing.
	Coliform count	Not to exceed 10 per gram.
	Drug residues	No positive results on drug residue detection methods which have been found to be acceptable for use with condensed whey.
DRY WHEY, EXTRA GRADE	Bacterial limit	Not to exceed 30,000 per gram.
	Coliform count	Not to exceed 10 per gram.
	Milkfat content	Not to exceed 1.5%.
	Moisture content	Not to exceed 5.0%.
	Scorched particle content	Not to exceed 15.0%.
DRY WHEY, DRY WHEY PRODUCTS	Bacterial limit	Not to exceed 50,000 per gram.
	Coliform count	Not to exceed 10 per gram.
	Butterfat	Not more than 1.50%.
	Moisture	Not more than 5%.
	Drug residues	No positive results on drug residue detection methods which have been found to be acceptable for use with dry whey and dry whey products.
DRY BUTTERMILK AND DRY BUTTERMILK PRODUCTS, EXTRA GRADE	Butterfat	4.5% min.
	Moisture	4.0% max.
	Titratable acidity	0.10-0.18%.
	Solubility index	1.25 ml spray process; 15.0 roller process.
	Bacterial limit	Not to exceed 20,000 per gram.
	Coliform count	Not to exceed 10 per gram.

	Scorched particles disc B	15.0 mg spray process; 22.5 mg roller process.
	Drug residues	No positive results on drug residue detection methods which have been found to be acceptable for use with dry buttermilk and dry buttermilk products.
DRY BUTTERMILK AND DRY BUTTERMILK PRODUCTS, STANDARD GRADE	Butterfat	4.5% min.
	Moisture	5.0% max.
	Titratable acidity	0.10-0.20%.
	Solubility index	2.0 ml spray process; 15.0 roller process.
	Bacterial limit	Not to exceed 75,000 per gram.
	Coliform count	Not to exceed 10 per gram.
	Scorched particles disc B	22.5 mg spray process; 32.5 mg roller process.
	Drug residues	No positive results on drug residue detection methods which have been found to be acceptable for use with dry buttermilk and dry buttermilk products.
BUTTER, WHIPPED BUTTER	Percent butterfat	Not less than 80%.
	Temperature	Maintained at a temperature of 45°F (7°C) or less, when in storage.
	Proteolytic count	Not more than 50 per gram.
	Yeast and mold	Not more than 10 per gram.
	Coliform count	Not more than 10 per gram.
	Keeping quality	Satisfactory after 7 days at 70°F (21°C).

PASTEURIZED MILK, CREAM, FLUID DAIRY PRODUCTS FOR FROZEN DESSERTS	Bacterial limit	Not to exceed 20,000 per ml.
	Coliform count	Not to exceed 10 per gram. Provided, that in the case of bulk milk transport tank shipments shall not exceed 100 per ml.
	Storage temp	No higher than 45°F (7°C).
FROZEN DESSERT MIX	Bacterial limit	30,000 per ml.
	Coliform count	Not to exceed 10 per gram. Provided, that in the case of bulk milk transport tank shipments shall not exceed 100 per ml.
	Storage temp	No higher than 45°F (7°C). (Sterile or aseptic mix has no storage temperature requirement.)
FROZEN DESSERTS	Bacterial limit	30,000 per ml.
	Coliform count	Not to exceed 10 per ml (20 per gram for chocolate, fruit, nuts, or other bulky flavored frozen desserts).
	Storage temp	No higher than 32°F (0°C).
	Butterfat	Per standards listed in 21 C.F.R. 135.
STERILIZED OR ASEPTIC PRODUCTS	Bacterial limit	Refer to 21 CFR 113.3(e)(1).
	Temperature	None.
	Yeast and mold	No viable yeast or mold spores.
	Drug residues	No positive results on drug residue detection methods which have been found to be acceptable for use with aseptically processed milk and milk products.

FOR DAIRY FARMS AND
DAIRY PLANTS;
RECIRCULATED COOLING
WATER (SWEET WATER);
GLYCOL FOR COOLING

as MPN or equivalent
method less than 1 per
100 ml.

CONDENSATE RECOVERY WATER (COW WATER)	Total plate count	Not to exceed 500 per ml.
	Chemical oxygen demand	Not to exceed 12 mg per L.
	Turbidity	Not to exceed 5 units.

History: 2001, Act 267, Eff. Feb. 8, 2002;—Am. 2008, Act 147, Eff. June 27, 2008.

Compiler's note: In the section of the table entitled "PASTEURIZED CONDENSED MILK AND CONDENSED SKIM MILK", the first instance of "145°" evidently should read "145°F".

288.631 Testing frequency for dairy products; testing dairy farm water supplies.

Sec. 71. (1) All dairy products shall be tested at a minimum frequency of 4 out of every 6 months, or as determined by the director.

(2) Except when the water supply is required to be tested annually as provided in section 130(9)(h), well water samples for dairy farms shall be tested a minimum of once every 3 years and found to be of safe and satisfactory quality.

(3) Water supplies for dairy plants shall be tested a minimum of once every 6 months.

(4) Recirculated water or recirculated cooling medium shall be tested a minimum of once every 6 months.

(5) Condensate recovery water shall be used only in applications that conform to requirements and procedures accepted by the United States food and drug administration or the director.

History: 2001, Act 267, Eff. Feb. 8, 2002.